arXiv:1612.08056v1 [gr-qc] 23 Dec 2016

Reentrant phase transitions of higher-dimensional AdS black holes in dRGT massive gravity

De-Cheng Zou* and Ruihong Yue†

College of Physical Science and Technology,

Yangzhou University, Yangzhou 225009, China

Ming Zhang‡

Faculty of Science, Xi'an Aeronautical University, Xi'an 710077,China

(Dated: December 26, 2016)

We study the $P-V$ criticality and phase transition in the extended phase space of AdS black holes in higher-dimensional dRGT massive gravity, treating the cosmological constant as pressure and the corresponding conjugate quantity is interpreted as thermodynamic volume. Besides the usual small/large black hole phase transitions, the interesting thermodynamic phenomena of reentrant phase transitions are observed for all $d \geq 6$ dimensions when the coupling coefficients $c_i m^2$ of massive potential satisfy some certain conditions.

PACS numbers: 04.50.Kd, 04.70.Dy, 04.50.Gh

I. INTRODUCTION

The Einstein's general relativity (GR) is a relativistic theory of gravity where the graviton is a massless spin-2 particle [1–3]. GR is the current description of gravitation in modern physics and has significantly astrophysical implications. Nevertheless, whether there exist a consistent extension of GR by a mass term is a basic challenge of classical field theory, since the open questions such as the old cosmological constant problem and the origin of the late-time acceleration of the Universe remain behind the puzzles at the interface between gravity/cosmology and particle physics. In general, by adding generic mass terms for the gravitons on the given background usually brings various instabilities for the gravitational theories, sometimes on the nonlinear level. A new nonlinear massive gravity theory was proposed by de Rham, Gabadadze and Tolley (dRGT) [4–6],

*Electronic address: dczou@yzu.edu.cn

†Electronic address: rhyue@yzu.edu.cn

‡Electronic address: shakellar@126.com

where the Boulware-Deser ghost [7] was eliminated by introducing higher order interaction terms in the action. Until now, a nontrivial black hole solution with a Ricci flat horizon in four-dimensional dRGT massive gravity has been constructed by Vegh [8, 9]. The spherically symmetric solutions for dRGT were also addressed in [10, 11], the corresponding charged black hole solution was found in [12], and its bi-gravity extension was found in [13]. Moreover, the charged AdS black hole solution in the higher-dimensional dRGT massive gravity, and its corresponding thermodynamics and phase structure in the grand canonical and canonical ensembles were also presented in [14].

Recently, the study of thermodynamics in AdS black holes has been generalized to the extended phase space, where the cosmological constant is regarded as a variable and also identified with thermodynamic pressure [15, 16]

$$P = -\frac{\Lambda}{8\pi} = \frac{(d-1)(d-2)}{16\pi l^2} \tag{1}$$

in the geometric units $G_N = \hbar = c = k = 1$. Here d stands for the spacetime dimensions and l denotes the AdS radius. In this case, the variation of the cosmological constant is included in the first law of black hole thermodynamics, which ensures the consistency between the first law of black hole thermodynamics and the Smarr formula. In [17], it was pointed out that the extended phase space can be interpreted as an RG-flow in the space of field theories, where isotherm curves codify how the number of degrees of freedom N (or the central charge c) runs with the energy scale. Moreover, the variation of cosmological constant could be corresponded to variation of number of the colors in Yang-Mills theory residing on the boundary spacetime [18, 19]. In the extended phase space, the charged AdS black hole admits a more direct and precise coincidence between the first order small/large black holes (SBH/LBH) phase transition and the Van der Waals liquid-gas phase transition, and both systems share the same critical exponents near the critical point [20]. As a result, the analogy between the charged AdS black hole and the Van der Waals system becomes more complete. More discussions were found as well in [21–42].

In this direction, some investigations for thermodynamics of AdS black holes in the dRGT massive gravity have been generalized to the extended phase space [43–47]. In these models, it revealed the existence of first-order SBH/LBH phase transition reminiscent of the liquid/gas transition of the Van der Waals fluid. On the other hand, the four-dimensional Born-Infeld-AdS [21], and higher-dimensional singly spinning AdS black holes [48] have been demonstrated the peculiar behavior of intermediate/small/large black hole phase transitions reminiscent of reentrant phase transitions (RPTs) observed for multicomponent fluid systems, ferroelectrics, gels, liquid crystals, and binary gases, e.g.,[49]. A system undergoes an RPT if a monotonic variation of

any thermodynamic quantity results in two (or more) phase transitions such that the final state is macroscopically similar to the initial state. In this paper, we will concentrate on the critical behaviors of higher-dimensional AdS black holes in the dRGT massive gravity, and also report the finding of interesting reentrant phase transition.

This paper is organized as follows. In Sect. II, we review the thermodynamics of massive gravity black holes in the extended phase space. In Sect. III, we study the critical behavior of higher-dimensional AdS black hole in context of $P-V$ criticality and phase diagrams. We end the paper with closing remarks in Sect. IV.

II. THERMODYNAMICS OF HIGHER-DIMENSIONAL ADS BLACK HOLE IN MASSIVE GRAVITY

We start with the action of higher-dimensional massive gravity in presence of a negative cosmological constant

$$\mathcal{I}=\frac{1}{16\pi}\int d^dx\sqrt{-g}\left[R+\Lambda+m^2\sum_{i=1}^{4}c_i\mathcal{U}_i(g,f)\right], \tag{2}$$

where the last four terms are the massive potential associate with graviton mass, c_i are constants and f is a fixed rank-2 symmetric tensor. Moreover, $\mathcal{U}_i$ are symmetric polynomials of the eigenvalues of the $d\times d$ matrix $\mathcal{K}^\mu_\nu\equiv\sqrt{g^{\mu\alpha}f_{\alpha\nu}}$

$$\begin{aligned}
\mathcal{U}_1 &= [\mathcal{K}],\\
\mathcal{U}_2 &= [\mathcal{K}]^2-[\mathcal{K}^2],\\
\mathcal{U}_3 &= [\mathcal{K}]^3-3[\mathcal{K}][\mathcal{K}^2]+2[\mathcal{K}^3],\\
\mathcal{U}_4 &= [\mathcal{K}]^4-6[\mathcal{K}^2][\mathcal{K}]^2+8[\mathcal{K}^3][\mathcal{K}]+3[\mathcal{K}^2]^2-6[\mathcal{K}^4].
\end{aligned} \tag{3}$$

The square root in $\mathcal{K}$ is understood as the matrix square root, ie., $(\sqrt{A})^\mu{}_\nu(\sqrt{A})_\lambda{}^\nu=A^\mu{}_\lambda$, and the rectangular brackets denote traces $[\mathcal{K}]=\mathcal{K}^\mu{}_\mu$.

Consider the metric of d-dimensional spacetime in the following form

$$ds^2=-f(r)dt^2+\frac{1}{f(r)}dr^2+r^2h_{ij}dx^idx^j, \tag{4}$$

where $h_{ij}dx^idx^j$ is the line element for an Einstein space with constant curvature $(d-2)(d-3)k$. The constant k characterizes the geometric property of hypersurface, which takes values $k=0$ for flat, $k=-1$ for negative curvature and $k=1$ for positive curvature, respectively.

By using the reference metric

$$f_{\mu\nu} = diag(0, 0, c_0^2 h_{ij}), \tag{5}$$

the metric function $f(r)$ is obtained as [14]

$$\begin{aligned} f(r) &= k + c_0^2 c_2 m^2 + \frac{16\pi P}{(d-1)(d-2)} r^2 + \frac{c_0 c_1 m^2}{d-2} r - \frac{16\pi M}{(d-2) V_{d-2} r^{d-3}} \\ &+ \frac{(d-3) c_0^3 c_3 m^2}{r} + \frac{(d-3)(d-4) c_0^4 c_4 m^2}{r^2}. \end{aligned} \tag{6}$$

Here c_0 is a positive constant, V_{d-2} is the volume of space spanned by coordinates x^i, and M is the black hole mass. It is worth to note that the terms $c_3 m^2$ and $c_4 m^2$ only appear in the black hole solutions for $d \geq 5$ and $d \geq 6$, respectively [14].

In terms of the horizon radius r_+, the mass M, Hawking temperature T and entropy S of black holes can be written as

$$\begin{aligned} M &= \frac{(d-2) V_{d-2} r_+^{d-3}}{16\pi} \left[k + \frac{16\pi P}{(d-1)(d-2)} r_+^2 + \frac{c_0 c_1 m^2 r_+}{d-2} + c_0 c_2 m^2 + \frac{(d-3) c_0^3 c_3 m^2}{r_+} \right. \\ &\left. + \frac{(d-3)(d-4) c_0^4 c_4 m^2}{r_+^2} \right], \\ T &= \frac{f'(r_+)}{4\pi} = \frac{1}{4\pi r_+} \left[(d-3)k + \frac{16\pi P}{d-2} r_+^2 + c_0 c_1 m^2 r_+ + (d-3) c_0^2 c_2 m^2 \right. \\ &\left. + \frac{(d-3)(d-4) c_0^3 c_3 m^2}{r_+} + \frac{(d-3)(d-4)(d-5) c_0^4 c_4 m^2}{r_+^2} \right], \\ S &= \frac{V_{d-2}}{4} r_+^{d-2}. \end{aligned} \tag{7}$$

The black hole mass M can be considered as the enthalpy rather than the internal energy of the gravitational system. Moreover, the first law of black hole thermodynamics and Smarr relation are given by [43]

$$\begin{aligned} dM &= TdS + VdP + \frac{c_0 m^2 V_{d-2} r_+^{d-2}}{16\pi} dc_1 + \frac{(d-2) c_0^2 m^2 V_{d-2} r_+^{d-3}}{16\pi} dc_2 \\ &+ \frac{(d-2)(d-3) c_0^3 m^2 V_{d-2} r_+^{d-4}}{16\pi} dc_3 + \frac{(d-2)(d-3)(d-4) c_0^4 m^2 V_{d-2} r_+^{d-5}}{16\pi} dc_4, \end{aligned} \tag{8}$$

$$\begin{aligned} (d-3)M &= (d-2)TS - 2VP - \frac{c_0 c_1 m^2 V_{d-2}}{16\pi} r_+^{d-2} + \frac{(d-2)(d-3) c_0^3 c_3 m^2 V_{d-2}}{16\pi} r_+^{d-4} \\ &+ \frac{(d-2)(d-3)(d-4) c_0^4 c_4 m^2 V_{d-2} r_+^{d-5}}{8\pi}, \end{aligned} \tag{9}$$

where V denotes the thermodynamic volume and equals to $V = \frac{V_{d-2}}{d-1} r_+^{d-1}$.

III. CRITICAL BEHAVIORS OF HIGHER-DIMENSIONAL ADS BLACK HOLES

A. Equation of state

For further convenience, we would like to denote

$$\hat{T} = T - \frac{c_0 c_1 m^2}{4\pi}, \quad w_2 = -\frac{k + c_0^2 c_2 m^2}{8\pi},$$
$$w_3 = -\frac{c_0^3 c_3 m^2}{8\pi}, \quad w_4 = -\frac{c_0^4 c_4 m^2}{8\pi}, \tag{10}$$

Here $\hat{T}$ denotes the shifted temperature and can be negative according to the value of $c_0 c_1 m^2$. Then, the equation of state of the black hole can be obtained from Eq. (7)

$$P = \frac{d-2}{4r_+}\left[\hat{T} + \frac{2(d-3)w_2}{r_+} + \frac{2(d-3)(d-4)w_3}{r_+^2} + \frac{2(d-3)(d-4)(d-5)w_4}{r_+^3}\right]. \tag{11}$$

To compare with the Van der Waals fluid equation, we can translate the "geometric" equation of state to physical one by identifying the specific volume v of the fluid with the horizon radius of the black hole as $v = \frac{4r_+}{d-2}$. Evidently, the specific volume v is proportional to the horizon radius r_+, therefore we will just use the horizon radius in the equation of state for the black hole hereafter in this paper.

We know that the critical point occurs when P has an inflection point,

$$\frac{\partial P}{\partial r_+}\Big|_{\hat{T}=\hat{T}_c, r_+=r_c} = \frac{\partial^2 P}{\partial r_+^2}\Big|_{\hat{T}=\hat{T}_c, r_+=r_c} = 0, \tag{12}$$

where the subscript stands for the quantities at the critical point. The critical shifted temperature is obtained as

$$\hat{T}_c = -\frac{2(d-3)}{r_c}\left[2w_2 + \frac{3(d-4)w_3}{r_c} + \frac{4(d-4)(d-5)w_4}{r_c^2}\right], \tag{13}$$

and the equation for the critical horizon radius r_c is given by

$$6(d-4)(d-5)w_4 + 3(d-4)w_3 r_c + w_2 r_c^2 = 0. \tag{14}$$

One can easily find that in four-dimensional spacetime ($d = 4, w_3 = w_4 = 0$), the absence of positive solution of Eq. (14) indicates that no criticality can happen [43].

We further discuss the critical behaviors of higher-dimensional ($d \geq 5$) AdS black hole when $w_2 \neq 0$ and $w_3 \neq 0$. When setting $w_4 = 0$, one have

$$r_c = -\frac{3(d-4)w_3}{w_2}, \quad \hat{T}_c = \frac{2(d-3)w_2^2}{3(d-4)w_3},$$
$$P_c = -\frac{(d-2)(d-3)w_2^3}{54(d-4)^2 w_3^2}. \tag{15}$$

Note that the critical behavior occurs only when $w_2 < 0$ and $w_3 > 0$. We can easily find an interesting relation among the critical pressure P_c, temperature $\hat{T}_c$ and horizon radius r_c

$$\frac{P_c r_c}{\hat{T}_c} = \frac{d-2}{12}. \tag{16}$$

For $d = 5$, the expressions (Eqs. (15)(16)) reduce to that described in [43]

With regard to the case of $w_4 \neq 0$, which only appears for $d \geq 6$, the direct solution of Eq. (14) reads as

$$\begin{aligned} r_{c1,2} \;=\; & \frac{1}{2w_2}\Big[\pm\sqrt{9(d-4)^2 w_3^2 - 24(d-4)(d-5)w_2 w_4} \\ & -3(d-4)w_3\Big] \equiv \frac{\chi_\pm}{2w_2} \end{aligned} \tag{17}$$

when $3(d-4)w_3^2 \geq 8(d-5)w_2 w_4$. In this case, r_{c1} and r_{c2} correspond to the " $-$ " and " $+$ " branches, respectively. The condition of $r_{c1,2} > 0$ crucially depends on the dimension of spacetime and values of w_2, w_3 and w_4.

For $w_2 < 0$, the positivity of solution r_{c1} leads to $w_3 < 0$ and $w_4 > 0$ or $w_3 > 0$. In order that r_{c2} be positive, it requires an additional constraint $w_3 > 0$ and $w_4 < 0$. Moreover, by substituting the solutions $r_{c1,2}$ (Eq. (17)) into Eqs. (11)(13), we obtain

$$\hat{T}_{c1,2} = \frac{8(d-3)(d-4)w_2^2\left[16(d-5)w_2 w_4 + 3w_3\chi_\pm\right]}{\chi_\pm^3}, \tag{18}$$

$$P_{c1,2} = \frac{\pm 4(d-2)(d-3)(d-4)w_2^3\left[6(d-5)w_2 w_4 + w_3\chi_\pm\right]}{\chi_\pm^4}. \tag{19}$$

Due to the shifted temperature $\hat{T}_{c1,2}$ can be negative, here we only evaluate the results of $P_{c1,2}$. When $P_{c1} > 0$, we shall keep $w_3 < 0$ and $w_4 > 0$ or $w_3 > 0$. On the other hand, taking $w_3 > 0$, $w_4 < 0$ and $\frac{3(d-4)w_3^2}{8(d-5)w_4} < w_2 < \frac{3(d-4)w_3^2}{9(d-5)w_4}$ lead to $P_{c2} > 0$. As a result, two critical points ($r_{c1,2} > 0$ and $P_{c1,2} > 0$) will appear in the range of $\frac{3(d-4)w_3^2}{8(d-5)w_4} < w_2 < \frac{3(d-4)w_3^2}{9(d-5)w_4}$ with $w_3 > 0$ and $w_4 < 0$.

In case of $w_2 > 0$, there is only one critical point ($r_{c1} > 0$ and $P_{c1} > 0$) for $w_3 < 0$, $w_4 > 0$ and $w_2 < \frac{(d-4)w_3^2}{3(d-5)w_4}$. We summarize the critical points in the Table I. The corresponding $P - r_+$ diagrams for $d = 6$ are displayed in Figs.1 and 2.

To study the possible phase transitions in the system, let us now turn to the expression for the Gibbs free energy.

TABLE I: The behaviors of critical points for different values of coupling constants in higher-dimensional spacetime.

parameters	$w_2 < 0$			$w_2 > 0$
w_3	$w_3 > 0$		$w_3 < 0$	$w_3 < 0$
w_4	$w_4 \geq 0$	$w_4 < 0$	$w_4 \geq 0$	$w_4 > 0$
Number of critical point	one	two	one	one



(a)$w_4 = 0.1$ (b)$w_4 = -0.7$

FIG. 1: **The $P - r_+$ diagrams of six dimensional AdS black holes for $w_2 = -1$ and $w_3 = 1$.** (a) The upper dashed line correspond to the idea gas phase behavior for $\hat{T} > \hat{T}_c$. The critical temperature case $\hat{T} = \hat{T}_c$ is denoted by the solid line. The line below is with temperatures smaller than the critical temperature. We have $\hat{T}_c = 0.9788$ in (a). In (b), we have now two critical points at positive pressure. The upper one has higher radius, temperature, and mass. We have $\hat{T}_{c1} = 1.2772$ and $\hat{T}_{c2} = 1.1718$ in (b).

B. Gibbs free energy

The behavior of free energy G is important to determine the thermodynamic phase transition. The free energy G obeys the thermodynamic relation

$$G = M - TS = -V_{d-2} r_+^{d-3} \left[\frac{P r_+^2}{(d-1)(d-2)} + \frac{w_2}{2} + \frac{(d-3)w_3}{r_+} + \frac{3(d-3)(d-4)w_4}{2r_+^2} \right]. \tag{20}$$

Here r_+ is understood as a function of pressure and temperature, $r_+ = r_+(P, \hat{T})$, via equation of state [Eq. (11)].

In the range of $\frac{3(d-4)w_3^2}{8(d-5)w_4} < w_2 < \frac{3(d-4)w_3^2}{9(d-5)w_4}$ with $w_3 > 0$ and $w_4 < 0$, the behavior of G in the six dimensional spacetime is depicted in Fig.3(a). We has one physical (with positive pressure) critical point and the corresponding first order SBH/LBH phase transition. This phase transition occurs for $\hat{T} < \hat{T}_{c1}$ and terminates at $\hat{T} = \hat{T}_t$. There also exists a certain range of temperatures,



(a)$w_2 = -1$, $w_3 = -1$ and $w_4 = 0.6$ (b)$w_2 = 1$, $w_3 = -1$ and $w_4 = 0.5$

FIG. 2: **The $P - r_+$ diagrams of six dimensional AdS black holes.** The upper dashed line correspond to the idea gas phase behavior for $\hat{T} > \hat{T}_c$. The critical temperature case $\hat{T} = \hat{T}_c$ is denoted by the solid line. The lines below are with temperatures smaller than the critical temperature. We have $\hat{T}_c = 19.2290$ in (a) and $\hat{T}_c = 1.1547$ in (b).

$\hat{T} \in (\hat{T}_t, \hat{T}_z)$, for which the global minimum of G is discontinuous, see Fig.3(b). In this range of temperatures, two separate branches of intermediate size and small size black holes co-exist. They are separated by a finite jump in G, which is so-called "zeroth-order phase transition". This phenomenon is also seen in superfluidity and superconductivity [50].



(a) (b)

FIG. 3: **Gibbs free energy of six dimensional AdS black holes for $w_2 = -1$ and $w_3 = 1$ and $w_4 = -0.7$.** In this range one has two critical points with positive pressure.

This novel situation can be also clearly illustrated in the $P - \hat{T}$ diagrams in Fig.4. There is the expected SBH/LBH line of coexistence, which ends in a critical point $(\hat{T}_{c1}, P_{c1})$ and terminates at $(\hat{T}_t, P_t)$. Especially, a "triple point" between the small, intermediate, and large black holes appears in the point $(\hat{T}_t, P_t)$. For $\hat{T} \in (\hat{T}_t, \hat{T}_z)$, there is a new IBH/SBH line of coexistence that terminates

in another critical point $(\hat{T}_z, P_z)$. The range for the RPT is quite narrow and must be determined numerically. Taking $w_2 = -1$, $w_3 = 1$, $w_4 = -0.7$ and $d = 6$, we obtain

$$(\hat{T}_t, \hat{T}_z, \hat{T}_{c1}) \approx (1.22194, 1.22459, 1.27568),$$
$$(P_t, P_z, P_{c1}) \approx (0.08615, 0.08747, 0.09860). \tag{21}$$



(a) (b)

FIG. 4: $P - \hat{T}$ **diagram of six dimensional AdS black holes for** $w_2 = -1$**,** $w_3 = 1$ **and** $w_4 = -0.7$. The coexistence line of the first order SBH/LBH phase transition is depicted by a thick solid line, which initiates from the critical point $(P_{c1}, \hat{T}_{c1})$ and terminates at $(P_t, \hat{T}_t)$. The solid line in the inset indicates the coexistence line of small and intermediate black holes, separated by a finite gap in G, indicating the RPT. It commences from $(P_z, \hat{T}_z)$ and terminates at $(P_t, \hat{T}_t)$.

In Fig. 5, we also plot the behaviors of Gibbs free energy of six dimensional AdS black hole for three other cases, showed in Table I. One can see that the G surface demonstrates the characteristic "swallow tail" behavior, which indicates that there is a Van der Waals like first order SBH/LBH phase transition when $P < P_c$ in the corresponding system. Moreover, the corresponding $P - T$ diagram (not shown) is reminiscent of what was observed for charged black holes in [20] and is analogous to the Van der Waals $P - T$ diagram in each case.

IV. CLOSING REMARKS

In this paper we have studied the thermodynamic behaviors of higher-dimensional AdS black hole in the dRGT massive gravity. We discussed this issue in the extended phase space where the cosmological constant appears as the pressure of the thermodynamic system and its conjugate quantity is the thermodynamic volume of the black holes. Following the standard thermodynamic techniques, we have written out the equations of state and examined the phase structures. When



(a)$w_2 = -1$, $w_3 = 1$ and $w_4 = 0.1$



(b)$w_2 = -1$, $w_3 = -1$ and $w_4 = 0.6$



(c)$w_2 = 1$, $w_3 = -1$ and $w_4 = 0.5$

FIG. 5: **Gibbs free energy for $d = 6$.** The behavior of the Gibbs free energy is depicted as a function of temperature for fixed pressure. There is one critical point and the corresponding first order SBH/LBH phase transition for $\hat{T} < \hat{T}_c$.

the coupling coefficients of massive potential satisfy $\frac{3(d-4)w_3^2}{8(d-5)w_4} < w_2 < \frac{3(d-4)w_3^2}{9(d-5)w_4}$ with $w_3 > 0$ and $w_4 < 0$, we find that a monotonic lowering of the temperature yields a large-small-large black hole transition, where we refer to the latter "large" state as an intermediate black hole (IBH), which is reminiscent of reentrant phase transitions. Moreover, this process is also accompanied by a discontinuity in the global minimum of the Gibbs free energy, referred to as a zeroth-order phase transition. For three other cases in Table I, the usual Van der Waals like small/large black hole phase transition occur when coupling coefficients of massive potential adopt some proper values in the higher-dimensional spacetime.

In addition, the charged black hole [51], Born-Infeld black hole [52], and black hole in the Maxwell and Yang-Mills fields [53] have been recently constructed in Gauss-Bonnet massive gravity. It has also showed the existence of Van der Waals like first order SBH/LBH phase transition in these models. It would be interesting to extend our discussion to these black hole solutions and find out whether the reentrant phase transition can appear.

Acknowledgments

The work is supported by the National Natural Science Foundation of China (Grant No.11605152, No.11275099, and No.11647050), and Natural Science Foundation of Jiangsu Province under Grant No.BK20160452.

[1] S. N. Gupta, "Gravitation and Electromagnetism," Phys. Rev. **96**, 1683 (1954).

[2] S. Weinberg, "Photons and gravitons in perturbation theory: Derivation of Maxwell's and Einstein's equations," Phys. Rev. **138**, B988 (1965).

[3] R. P. Feynman, F. B. Morinigo, W. G. Wagner and B. Hatfield, "Feynman lectures on gravitation," Reading, USA: Addison-Wesley (1995) 232 p. (The advanced book program)

[4] C. de Rham and G. Gabadadze, "Generalization of the Fierz-Pauli Action," Phys. Rev. D **82**, 044020 (2010) [arXiv:1007.0443 [hep-th]].

[5] C. de Rham, G. Gabadadze and A. J. Tolley, "Resummation of Massive Gravity," Phys. Rev. Lett. **106**, 231101 (2011) [arXiv:1011.1232 [hep-th]].

[6] K. Hinterbichler, "Theoretical Aspects of Massive Gravity," Rev. Mod. Phys. **84**, 671 (2012) [arXiv:1105.3735 [hep-th]].

[7] D. G. Boulware and S. Deser, "Can gravitation have a finite range?," Phys. Rev. D **6**, 3368 (1972).

[8] D. Vegh, "Holography without translational symmetry," arXiv:1301.0537 [hep-th].

[9] A. Adams, D. A. Roberts and O. Saremi, "Hawking-Page transition in holographic massive gravity," Phys. Rev. D **91**, no. 4, 046003 (2015) [arXiv:1408.6560 [hep-th]].

[10] T. M. Nieuwenhuizen, "Exact Schwarzschild-de Sitter black holes in a family of massive gravity models," Phys. Rev. D **84**, 024038 (2011) [arXiv:1103.5912 [gr-qc]].

[11] R. Brito, V. Cardoso and P. Pani, "Black holes with massive graviton hair," Phys. Rev. D **88**, 064006 (2013) [arXiv:1309.0818 [gr-qc]].

[12] L. Berezhiani, G. Chkareuli, C. de Rham, G. Gabadadze and A. J. Tolley, "On Black Holes in Massive Gravity," Phys. Rev. D **85**, 044024 (2012) [arXiv:1111.3613 [hep-th]].

[13] E. Babichev and A. Fabbri, "A class of charged black hole solutions in massive (bi)gravity," JHEP **1407**, 016 (2014) [arXiv:1405.0581 [gr-qc]].

[14] R. G. Cai, Y. P. Hu, Q. Y. Pan and Y. L. Zhang, "Thermodynamics of Black Holes in Massive Gravity," Phys. Rev. D **91**, 024032 (2015) [arXiv:1409.2369 [hep-th]].

[15] B. P. Dolan, "Pressure and volume in the first law of black hole thermodynamics," Class. Quant. Grav. **28**, 235017 (2011) [arXiv:1106.6260 [gr-qc]].

[16] B. P. Dolan, "The cosmological constant and the black hole equation of state," Class. Quant. Grav. **28**, 125020 (2011) [arXiv:1008.5023 [gr-qc]].

[17] E. Caceres, P. H. Nguyen and J. F. Pedraza, "Holographic entanglement entropy and the extended

phase structure of STU black holes," JHEP **1509**, 184 (2015) [arXiv:1507.06069 [hep-th]].

[18] J. D. E. Creighton and R. B. Mann, "Quasilocal thermodynamics of dilaton gravity coupled to gauge fields," Phys. Rev. D **52**, 4569 (1995) [gr-qc/9505007].

[19] G. W. Gibbons, R. Kallosh and B. Kol, "Moduli, scalar charges, and the first law of black hole thermodynamics," Phys. Rev. Lett. **77**, 4992 (1996) [hep-th/9607108].

[20] D. Kubiznak and R. B. Mann, "P-V criticality of charged AdS black holes," JHEP **1207**, 033 (2012) [arXiv:1205.0559 [hep-th]].

[21] S. Gunasekaran, R. B. Mann and D. Kubiznak, "Extended phase space thermodynamics for charged and rotating black holes and Born-Infeld vacuum polarization," JHEP **1211**, 110 (2012) [arXiv:1208.6251 [hep-th]].

[22] S. H. Hendi and M. H. Vahidinia, "P-V criticality of higher dimensional black holes with nonlinear source," Phys. Rev. D **88**, 084045 (2013) [arXiv:1212.6128 [hep-th]].

[23] S. Chen, X. Liu, C. Liu and J. Jing, "$P-V$ criticality of AdS black hole in $f(R)$ gravity," Chin. Phys. Lett. **30**, 060401 (2013) [arXiv:1301.3234 [gr-qc]].

[24] R. G. Cai, L. M. Cao, L. Li and R. Q. Yang, "P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space," JHEP **1309**, 005 (2013) [arXiv:1306.6233 [gr-qc]].

[25] S. Dutta, A. Jain and R. Soni, "Dyonic Black Hole and Holography," JHEP **2013**, 60 (2013) [arXiv:1310.1748 [hep-th]].

[26] W. Xu, H. Xu and L. Zhao, "Gauss-Bonnet coupling constant as a free thermodynamical variable and the associated criticality," Eur. Phys. J. C **74**, 2970 (2014) [arXiv:1311.3053 [gr-qc]].

[27] D. C. Zou, S. J. Zhang and B. Wang, "Critical behavior of Born-Infeld AdS black holes in the extended phase space thermodynamics," Phys. Rev. D **89**, 044002 (2014) [arXiv:1311.7299 [hep-th]].

[28] H. H. Zhao, L. C. Zhang, M. S. Ma and R. Zhao, "P-V criticality of higher dimensional charged topological dilaton de Sitter black holes," Phys. Rev. D **90**, 064018 (2014).

[29] N. Altamirano, D. Kubiznak, R. B. Mann and Z. Sherkatghanad, "Thermodynamics of rotating black holes and black rings: phase transitions and thermodynamic volume," Galaxies **2**, 89 (2014) [arXiv:1401.2586 [hep-th]].

[30] J. -X. Mo and W. -B. Liu, "P-V Criticality of Topological Black Holes in Lovelock-Born-Infeld Gravity," Eur. Phys. J. C **74**, 2836 (2014) [arXiv:1401.0785 [gr-qc]].

[31] D. C. Zou, Y. Liu and B. Wang, "Critical behavior of charged Gauss-Bonnet AdS black holes in the grand canonical ensemble," Phys. Rev. D **90**, 044063 (2014) [arXiv:1404.5194 [hep-th]].

[32] H. Xu, W. Xu and L. Zhao, "Extended phase space thermodynamics for third order Lovelock black holes in diverse dimensions," Eur. Phys. J. C **74**, 3074 (2014) [arXiv:1405.4143 [gr-qc]].

[33] W. Xu and L. Zhao, "Critical phenomena of static charged AdS black holes in conformal gravity," Phys. Lett. B **736**, 214 (2014) [arXiv:1405.7665 [gr-qc]].

[34] M. H. Dehghani, S. Kamrani and A. Sheykhi, "$P-V$ criticality of charged dilatonic black holes," Phys. Rev. D **90**, 104020 (2014).

[35] C. O. Lee, "The extended thermodynamic properties of Taub-NUT/Bolt-AdS spaces," Phys. Lett. B **738**, 294 (2014) [arXiv:1408.2073 [hep-th]].

[36] J. L. Zhang, R. G. Cai and H. Yu, "Phase transition and thermodynamical geometry for Schwarzschild AdS black hole in $AdS_5 \times S^5$ spacetime," JHEP **1502**, 143 (2015) [arXiv:1409.5305 [hep-th]].

[37] M. Zhang, Z. Y. Yang, D. C. Zou, W. Xu and R. H. Yue, "$P - V$ criticality of AdS black hole in the Einstein-Maxwell-power-Yang-Mills gravity," Gen. Rel. Grav. **47**, 14 (2015) [arXiv:1412.1197 [hep-th]].

[38] M. Zhang, R. H. Yue and Z. Y. Yang, "Critical behaviors of black holes in an Einstein-Maxwell gravity with a conformal anomaly," Chin. Phys. Lett. **32**, 020401 (2015) [arXiv:1412.1201 [hep-th]].

[39] J. L. Zhang, R. G. Cai and H. Yu, "Phase transition and thermodynamical geometry of Reissner-Nordstrom-AdS black holes in extended phase space," Phys. Rev. D **91**, 044028 (2015) [arXiv:1502.01428 [hep-th]].

[40] Y. Liu, D. C. Zou and B. Wang, "Signature of the Van der Waals like small-large charged AdS black hole phase transition in quasinormal modes," JHEP **1409**, 179 (2014) [arXiv:1405.2644 [hep-th]].

[41] B. R. Majhi and S. Samanta, "P-V criticality of AdS black holes in a general framework," arXiv:1609.06224 [gr-qc].

[42] S. H. Hendi, B. Eslam Panah, S. Panahiyan and M. S. Talezadeh, "Geometrical thermodynamics and P-V criticality of black holes with power-law Maxwell field," arXiv:1612.00721 [hep-th].

[43] J. Xu, L. M. Cao and Y. P. Hu, "P-V criticality in the extended phase space of black holes in massive gravity," Phys. Rev. D **91**, 124033 (2015) [arXiv:1506.03578 [gr-qc]].

[44] S. H. Hendi, B. Eslam Panah and S. Panahiyan, "Einstein-Born-Infeld-Massive Gravity: adS-Black Hole Solutions and their Thermodynamical properties," JHEP **1511**, 157 (2015) [arXiv:1508.01311 [hep-th]].

[45] M. Zhang and W. B. Liu, "Coexistent physics of massive black holes in the phase transitions," arXiv:1610.03648 [gr-qc].

[46] S. G. Ghosh, L. Tannukij and P. Wongjun, "A class of black holes in dRGT massive gravity and their thermodynamical properties," Eur. Phys. J. C **76**, 3 (2016) [arXiv:1506.07119 [gr-qc]].

[47] P. Prasia and V. C. Kuriakose, "Quasi Normal Modes and P-V Criticallity for scalar perturbations in a class of dRGT massive gravity around Black Holes," Gen. Rel. Grav. **48**, 89 (2016) [arXiv:1606.01132 [gr-qc]].

[48] N. Altamirano, D. Kubiznak and R. B. Mann, "Reentrant Phase Transitions in Rotating AdS Black Holes," Phys. Rev. D **88**, 101502 (2013) [arXiv:1306.5756 [hep-th]].

[49] T. Narayanan and A. Kumar, "Reentrant phase transitions in multicomponent liquid mixtures," Physics Reports **249** (1994).

[50] V. P. Maslov, "Zeroth-order phase transitions," *Mathematical Notes* **76** 697-710 (2004)

[51] S. H. Hendi, S. Panahiyan and B. Eslam Panah, "Charged Black Hole Solutions in Gauss-Bonnet-Massive Gravity," JHEP **1601**, 129 (2016) [arXiv:1507.06563 [hep-th]].

[52] S. H. Hendi, G. Q. Li, J. X. Mo, S. Panahiyan and B. Eslam Panah, "New perspective for black

hole thermodynamics in Gauss-Bonnet-Born-Infeld massive gravity," Eur. Phys. J. C **76**, 10 (2016) [arXiv:1608.03148 [gr-qc]].

[53] K. Meng and J. Li, "Black hole solution of Gauss-Bonnet massive gravity coupled to Maxwell and Yang-Mills fields in five dimensions," Europhys. Lett. **116**, 10005 (2016).